Low-temperature Thermodynamics of a fermi liquid.

To calculate the specific heat, we simply note that the deading contribution comed from states close to the fermi surface with energy (2p-4/5) T.

As just discussed, at this order,  $N_{p}(T) = \frac{1}{e^{\beta(\xi_{pq}^{2} - \mu)} + 1}, \text{ and therefore}$ 

one may carry out the standard Sommerfeld expansion, leading to

$$\frac{C(V,T,N)}{V} = \frac{T^2}{2} \frac{Tn}{\epsilon_F}$$

where  $N = \frac{N}{V}$  and  $EF = \frac{P_F^2}{2m^*}$  N and  $P_F$  are related of usual:  $N = \frac{P_F^3}{3\pi^2}$ . Note again that  $P_F$ 

3 The unchanged compared to non-interacting es Chuttinger theorem), and the only

effect of interactions is that the electron mass is replaced by the effective mass m\*. Using above relations on man also write.

One may also write.  $\frac{C(T, V, N)}{c} = a T n^{1/3} m^*$ 

where a is a numerical constant. Therefore, the speaks heat is

Enhanced by a factor of  $\frac{m^*}{m}$ compared to the non-Interacting care. One may now calculate other grantities Such at entropy and the temperature dependence of the chamical potential. Entropy:  $T \frac{\partial x}{\partial T} = \frac{C}{V} = \alpha T n^{1/3} m^*$ where  $\Delta = \frac{SCT, V, N}{V}$ .  $\Rightarrow 3 = 2 n^{1/3} m^* T + \underbrace{f(N,V)}_{J_2(T=0)}$ When there are no particles i.e. when when N=0, we expect that A=0.

Forther,  $\Delta (T=0) = 0$  from the third loss of thermodynamics, = 2 + 2 + 2 = 0 = One may use the Maxwell relation.  $\frac{\partial \mu}{\partial T} = -\frac{\partial S}{\partial N} = -\frac{\partial (SN)}{\partial N}$ 

$$= - \alpha \left[ \frac{3}{1} \sqrt{2} \sqrt{3} + \sqrt{2} \sqrt{3} + \sqrt{3} \sqrt{3} \right]$$

Chemical potential:

 $=) \mu(T,n) = \mu(T=0) - \alpha T^{2} \left[ m^{*} \pi^{-2/3} + n^{1/3} \frac{\partial n}{\partial m^{*}} \right]$ 

We note that in addition to the standard contribution proportional to  $-72\pi^{2/3}m^*$ , here one gets

on additional term proportional to  $\frac{\partial m^*}{\partial n}$ . See Baym and Petrik for

on alternative derivation of C, Ja, K.

Compressibility at T=0

Compressibility K is given by  $k = -\frac{\partial V}{V \partial P}$ =  $\frac{1}{n^2} \frac{\partial N}{\partial \mu}$  where N is the density.

Using the integral equation for Npr.

[48 - 2938] = 3npe = 29NS

Note that 82pt = 855pp (npror-nor)
depends implicitly on 8p due to the

Voriation of Npr wir.t. 4.
Summing over all P, T

 $-\frac{6a}{2}\frac{9\epsilon^{6a}}{9u^{6a}} + \frac{9\epsilon^{6a}}{2u^{6a}} = \frac{6a}{2u^{6a}}\frac{9\epsilon^{6a}}{2u^{6a}} + \frac{6a}{2u^{6a}} + \frac{9\epsilon^{6a}}{2u^{6a}}$ 

As usual, the tacker of  $\frac{3n}{32}$  restricts both P and P' to Die at the Fermi surface  $\left[\frac{3n}{32} \sim -8(E-EF)\right]$ Let's consider the two terms on the right hand side separately.  $-\frac{2}{5}\frac{3n_{10}}{3E_{10}} \approx \mu = \frac{3}{5}\frac{2}{5} \approx 1000$   $= \frac{3}{5}$   $= \frac{3}$ 

where D(E) dendes density of states (we use D instead of N or n to wake distinction with particle number

make distriction with particle countries.

or graniparticle occuration).

The second term is  $\sum \frac{\partial n_{po}}{\partial x_{po}} \left[\sum_{p' \in Y} S_{pp'} S_{np' \in Y}\right]$ 

The second term is  $\frac{5}{2} \frac{\partial n_{po}}{\partial \epsilon_{p\sigma}} \left[ \frac{5}{2}, \frac{5}{60!}, \frac{5}{8} \frac{n_{p'e'}}{n_{p'e'}} \right]$ 

PP'00' 501 SNp'0'

Fermi surface

> for can be simplied The sum is expanded in angular 15 Sm! monentain modes in d=2, or Using Legendre Polynomials in d=3. First, oue utilized the spin-rotational Symmetry to write 2661 = 2662 + 266, QQ, where fs devotes the spin-symmetric and for the spin anti-symmetric component. 55 and 50 may then further be expanded in terms of angle & between band p': d=3:  $5\frac{s}{pp}=\frac{8}{9}5\frac{s}{9}$  Palcosa)

$$d=2$$
:  $S_{PP}^{S/Q} = \sum_{n=-\infty}^{\infty} S_{n}^{S/Q} e^{-n} d$   
Returning to the sum  $\sum_{n=0}^{\infty} S_{n}^{PP}^{S/Q} e^{-n} d$   
only the  $d=0$  symmetric part

oney the 
$$l=0$$
 symmetric part  
survives  $\sum_{pq} f_{pq} = V f_{q} = 0$   
 $= 0$ 

$$= \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} = \frac$$

Where	8 M	60	۵ <b>۱</b> ۱۱	60		
Combining	ere	ry tuin	g-,			-1
N8	2	D(22)	[ <	V — 43	52=0	[118]

$$8N = 18 = 18 = 0.8$$

where F2=0 = V f2=0 D(Ex)

everything,
$$= 2000 \times 1000 \times 1000$$

$$= 8\mu \times 1000 \times 1000$$

$$= 8\mu \times 1000 \times 1000$$

$$= 1 + F^{S}_{2} = 0$$

 $= \frac{18n}{\sqrt{20}}$ 

 $= \frac{u_5}{1} \frac{1 + E_c^{3=0}}{5 C E^{6}}.$ 

Compared to the non-interacting fermions

one trerelore obtains on additional factor of 1/1+FS=0. Also, recall that

the density of states DLEF) is

also enhanced by a factor of mit

compared to the non-interacting es.

(C) Spin susceptibility:

The spin-susceptibility is defined as X  $=\frac{\partial M}{\partial L}$  where M is magnetization

and h is the external magnetic

field. The derivation for X is almost identical to trat for K. Ept in the presence of magnetic field

h & given by 5 ba = 5m - 1 2 wba + 260, 8wba 8wba

The voriation of magnetization is  $\delta M = \sum_{i} \Delta_{i} \delta N \delta \Delta_{i} = \frac{\delta \alpha}{2} \frac{9 \epsilon^{\delta \alpha}}{9 N^{\delta \alpha} \Delta_{i}} \sum_{i} \frac{\delta \alpha}{2^{\delta \delta_{i}}} \delta U \delta_{i} \delta_{i}$ 

- 5 3062 8W

Again 
$$p, p' \in F.S.$$
 because of the factor  $\frac{\partial n}{\partial \varepsilon}$ . The sum  $\sum_{pr} \sigma \int_{\sigma \sigma'}^{pp'}$ 

$$= \int_{0}^{\alpha} q = 0 \quad \nabla \sigma'$$

$$= f_{0=0} \vee \sigma'$$

$$= 8m = -20 (2E) \vee f_{0=0} \otimes m$$

$$= \frac{8W}{1 + E_{0}} = \frac{8W}{1 + E_{0}}$$

Effective wass mx :

Remarkably, one may deduce a relation between  $\frac{M^*}{M}$  and Landon parameter  $F_{a=1}^s$ , if the system is Galilean invariant (e.g. a Single-component He-3, but not a

mixture of He-3 and He-4).

Let's denote the Hamiltonian in the lab frame of H, and consider a

frame that is moving at relocity it with the lab frame. The Hamiltonian H' and momentum P' in this frame are

and momentum  $\vec{P}'$  in this frame are  $H' = H - \vec{P} \cdot \vec{u} + \frac{1}{2} M u^2, \ \vec{P}' = \vec{P} - M \vec{u}'$ 

where P is the total momentum in the lab frame, and M is the total mass, which is frame invariant Checause the system is Galilean invariant).

From now on, we will call the frame.
moving wirt. lab as the primed frame. Consider adding a single fermion with momentum of to the system. He soon as it is added, it will get dressed by other fermions in the system, and become a grasipartile with womentum & and enersy EP =  $\frac{1}{2m}$ \* +  $\frac{5}{18}$  Spp. 8mp. (we ignore the spin index in this discussion us it doesn't matter). The total mass Of the system changes by M, the Mans of the bare fermion leg, mass of a single He-3 atom). = DE (change in enersy = in the lab frame) DP (change in momentum = p

 $\Delta M = m$ .

Using the above mentioned relation between relation between the two frames (i.e. the usual Gazilean transformation laws);

DE (Change in energy in the primes frame)

= DE - 3. DF + 1/2MW2 = EF - F. I + 1/2 mo<sup>2</sup>

DF (Change in momentum

in the primed frame)
$$= \Delta \vec{P} - (\Delta M) \vec{O} = \vec{P} - m \vec{O}$$

Therefore in the primed frame, the energy-momentum dispersion for a q.p. is

enersy-momentum dispersion for a q.p. 1.
$$\mathcal{E}_{\overline{p}-m\overline{q}} = \mathcal{E}_{\overline{p}-\overline{p}} \cdot \overline{\mathcal{I}} + \frac{1}{2} m u^{2}$$

or, equivalently,  $\mathcal{E} = \mathcal{E} + m\vec{x} - \vec{r} \cdot \vec{x} - \frac{1}{2} m \vec{u}^2.$ 

Lets expand the R-18-s. to OCU):  $\tilde{v} \cdot \tilde{q} - \tilde{v}m \cdot \frac{\tilde{q} \cdot 6}{\tilde{q} \cdot 6} + \tilde{q} \cdot 3 \approx \tilde{q} \cdot 3$ In the lab frame, there are no Excitations,  $\Rightarrow$   $\leq np = 0 \Rightarrow \leq p = \frac{p^2}{2m^*}$  $\tilde{v}_{0} = v_{0} = v$  $\overline{u} \cdot \overline{q} \left( L - \frac{m}{m} \right) + \overline{q} = 0$ On the other hand, in the primed frame, the distribution NP is a filled Permi sea contered at  $P_0 = -mu^*$ .

primed frame

to therefore, in this

frame frame, there are

exceltations a shown

in the figure

outile

controls

co Porticle - like excitations.

Can be calulated using Landou's ausatz! Ep = Ep + \( \frac{7}{p}, \frac{1}{9}p\_1 \) & \( \text{Np1} \) 8 Np1 = Np1+mu - Np1

The energy ust of these excitations

 $= \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial g_{0}}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}} \frac{\partial v}{\partial g_{0}} \cdot m\vec{u} = \frac{\partial v}{\partial g_{0}$ => E'p = Ep + \( \frac{2n}{m} \frac{n\dot{p}}{m\*} \) \( \frac{1}{m} \frac{2n}{m} \frac{n}{m} \frac{2}{m} \)

As usual, the factor of  $\frac{\partial n}{\partial s}$  restricts tue sum to the Ferni surface. Since

B1 15 a rector l'i.e. angular momentum

= 1), only for will contribute.

To find the exact pretactor, we use

/d(cose) f s=1 Po=160=(0-01) (cos(01)

 $= \frac{1}{3} f_{0=1}^{2} cos(\theta) \chi^{2}$   $= \frac{1}{3} f_{0=1}^{2} cos(\theta)$ 

$$=) \quad \mathcal{E}_{p} = \mathcal{E}_{p} - V \underbrace{f_{0=1}^{s}}_{\mathcal{T}_{m}} \mathcal{D}(E_{p}) \overrightarrow{p}. \overrightarrow{w}_{m}^{*}$$

$$= \varepsilon_{\rho} - \frac{F_{0=1}^{s}}{3} \overrightarrow{p} \cdot \overrightarrow{u} \frac{m}{m^{*}}$$

where 
$$FS_{2-1} = VS_{2-1} 2CEF)$$
  
Company with the ofterrate

expression for 
$$\mathcal{E}_{p}$$
 derived above,
$$\frac{m}{m} - 1 = -\frac{\mathcal{F}_{a=1}}{m}$$

$$\frac{m}{m*} - 1 = -\frac{1}{E_{s}} \frac{m}{m*}$$

$$\Rightarrow \frac{m}{m^*} = \frac{1}{1 + \frac{1}{2}}$$

$$\Rightarrow \frac{m^*}{m} = 1 + \frac{5}{3}$$